

K -Theory for Algebras of Operators on Banach Spaces

Niels Jakob Laustsen*

Department of Mathematics and Computer Science
Odense University, Campusvej 55, DK-5230 Odense M, Denmark
e-mail: laustsen@imada.ou.dk

Abstract

We prove that, for each pair (m, n) of non-negative integers, there is a Banach space $\mathfrak{X}$ for which $K_0(\mathcal{B}(\mathfrak{X})) \cong \mathbb{Z}^m$ and $K_1(\mathcal{B}(\mathfrak{X})) \cong \mathbb{Z}^n$. Along the way we compute the K -groups of all closed ideals of operators contained in the ideal of strictly singular operators, and we derive some results about the existence of splittings of certain short exact sequences.

1 Introduction and Notation

Throughout, all vector spaces and algebras are tacitly assumed to be complex. In addition, all Banach spaces are assumed to be infinite-dimensional. Banach spaces are denoted by $\mathfrak{X}$, $\mathfrak{Y}$, $\mathfrak{Z}$, and $\mathfrak{W}$. The term ‘operator’ means a continuous linear map. The Banach space of all operators from $\mathfrak{X}$ to $\mathfrak{Y}$ is denoted by $\mathcal{B}(\mathfrak{X}, \mathfrak{Y})$ (or $\mathcal{B}(\mathfrak{X})$ in the case where $\mathfrak{X} = \mathfrak{Y}$).

By an *ideal of operators* we understand an assignment $\mathcal{I}$ which associates to each pair $(\mathfrak{X}, \mathfrak{Y})$ of Banach spaces a subspace $\mathcal{I}(\mathfrak{X}, \mathfrak{Y})$ of $\mathcal{B}(\mathfrak{X}, \mathfrak{Y})$ satisfying

$$TSR \in \mathcal{I}(\mathfrak{W}, \mathfrak{Z}) \text{ whenever } R \in \mathcal{B}(\mathfrak{W}, \mathfrak{X}), S \in \mathcal{I}(\mathfrak{X}, \mathfrak{Y}), \text{ and } T \in \mathcal{B}(\mathfrak{Y}, \mathfrak{Z}).$$

In the case where $\mathfrak{X} = \mathfrak{Y}$, we write $\mathcal{I}(\mathfrak{X})$ instead of $\mathcal{I}(\mathfrak{X}, \mathfrak{X})$. An ideal of operators $\mathcal{I}$ is said to be *closed* provided that $\mathcal{I}(\mathfrak{X}, \mathfrak{Y})$ is a closed subspace of $\mathcal{B}(\mathfrak{X}, \mathfrak{Y})$ for all Banach spaces $\mathfrak{X}$ and $\mathfrak{Y}$, and $\mathcal{I}$ is said to be *non-zero* if $\mathcal{I}(\mathfrak{X}, \mathfrak{Y}) \neq \{0\}$ for all (non-zero) Banach spaces $\mathfrak{X}$ and $\mathfrak{Y}$. Let $\mathcal{I}_1$ and $\mathcal{I}_2$ be two ideals of operators. We say that $\mathcal{I}_1$ is *contained* in $\mathcal{I}_2$ provided that $\mathcal{I}_1(\mathfrak{X}, \mathfrak{Y}) \subseteq \mathcal{I}_2(\mathfrak{X}, \mathfrak{Y})$ for all Banach spaces $\mathfrak{X}$ and $\mathfrak{Y}$. Note that an ideal of operators is non-zero if and only if it contains the ideal $\mathcal{F}$ of finite-rank operators. The *rank* of a finite-rank operator A is denoted by $\text{rk } A$, i.e., $\text{rk } A = \dim(\text{im } A)$.

Recall that an operator $S : \mathfrak{X} \rightarrow \mathfrak{Y}$ is said to be *strictly singular* provided that, whenever $\tilde{\mathfrak{X}}$ is a subspace of $\mathfrak{X}$ on which S is bounded below, then $\tilde{\mathfrak{X}}$ is finite-dimensional

*Supported by The Danish Natural Science Research Council.

(cf. [Ka1]). The collection $\mathcal{S}$ of strictly singular operators forms a closed ideal of operators containing the closed ideal of compact operators.

Let $n \in \mathbb{N}$. We denote the direct sum of the Banach spaces $\mathfrak{X}_1, \dots, \mathfrak{X}_n$ by $\bigoplus_{j=1}^n \mathfrak{X}_j$ (or $\mathfrak{X}^n$ in the case where $\mathfrak{X}_1 = \dots = \mathfrak{X}_n = \mathfrak{X}$). This is a Banach space under coordinatewise operations and the maximum norm.

For operators $T_j \in \mathcal{B}(\mathfrak{X}_j, \mathfrak{Y}_j)$ ($j \in \{1, \dots, n\}$), we define

$$T_1 \oplus \dots \oplus T_n : (x_1, \dots, x_n) \longmapsto (T_1 x_1, \dots, T_n x_n), \quad \bigoplus_{j=1}^n \mathfrak{X}_j \longrightarrow \bigoplus_{j=1}^n \mathfrak{Y}_j.$$

This is clearly a linear and continuous map.

Similar notation and conventions apply to direct sums of (Banach) algebras and groups.

Let $\mathcal{A}$ be an algebra, and let $m, n \in \mathbb{N}$. We denote the vector space of $(m \times n)$ -matrices over $\mathcal{A}$ by $M_{m,n}(\mathcal{A})$ (or $M_n(\mathcal{A})$ in the case where $m = n$). An algebra homomorphism $\varphi : \mathcal{A} \rightarrow \mathcal{B}$ induces an algebra homomorphism $\varphi_n : M_n(\mathcal{A}) \rightarrow M_n(\mathcal{B})$ by the definition

$$\varphi_n \left((A_{kl})_{k,l=1}^n \right) := (\varphi(A_{kl}))_{k,l=1}^n.$$

Suppose that $\mathcal{A}$ is unital. We write I for the identity of $\mathcal{A}$, and the group of invertible elements in $\mathcal{A}$ is denoted by $\text{Inv}(\mathcal{A})$. The identity of $M_n(\mathcal{A})$ is denoted by $I^{(n)}$, and we let $\text{Inv}_n(\mathcal{A}) := \text{Inv}(M_n(\mathcal{A}))$, the invertible group in $M_n(\mathcal{A})$.

For a non-unital algebra $\mathcal{A}$, we denote $\mathcal{A}$ with an identity adjoined by $\mathcal{A}^\#$. Suppose that $\mathcal{B}$ is a unital algebra containing $\mathcal{A}$ as a subalgebra. Then we identify $\mathcal{A}^\#$ with the subalgebra $\mathcal{A} + \mathbb{C}I$ of $\mathcal{B}$.

Now suppose that $\mathcal{A}$ is a unital Banach algebra. To make $M_{m,n}(\mathcal{A})$ a Banach space (or a Banach algebra in the case where $m = n$), we consider it as a closed subspace (subalgebra) of $\mathcal{B}(\mathcal{A}^n, \mathcal{A}^m)$ in the obvious way. The norm on $M_{m,n}(\mathcal{A})$ thus obtained satisfies

$$\begin{aligned} \max \{ \|A_{kl}\| \mid k \in \{1, \dots, m\}, l \in \{1, \dots, n\} \} &\leq \left\| (A_{kl})_{k,l=1}^{m,n} \right\| \\ &\leq \max \left\{ \sum_{l=1}^n \|A_{kl}\| \mid k \in \{1, \dots, m\} \right\}. \end{aligned}$$

Let $\mathfrak{X}_1, \dots, \mathfrak{X}_n$ and $\mathfrak{Y}_1, \dots, \mathfrak{Y}_m$ be Banach spaces. There is a standard bijective correspondence between operators $T \in \mathcal{B} \left(\bigoplus_{j=1}^n \mathfrak{X}_j, \bigoplus_{j=1}^m \mathfrak{Y}_j \right)$ and $(m \times n)$ -matrices $(T_{kl})_{k,l=1}^{m,n}$ with $T_{kl} \in \mathcal{B}(\mathfrak{X}_l, \mathfrak{Y}_k)$, and for every ideal of operators $\mathcal{I}$ we have:

$$T \in \mathcal{I} \left(\bigoplus_{j=1}^n \mathfrak{X}_j, \bigoplus_{j=1}^m \mathfrak{Y}_j \right) \text{ if and only if } T_{kl} \in \mathcal{I}(\mathfrak{X}_l, \mathfrak{Y}_k) \text{ for all } k, l. \quad (1.1)$$

In the sequel we *identify* the operator T with its matrix representation $(T_{kl})_{k,l=1}^{m,n}$. In particular, $\mathcal{B}(\mathfrak{X}^n, \mathfrak{X}^m)$ is identified with $M_{m,n}(\mathcal{B}(\mathfrak{X}))$ for each Banach space $\mathfrak{X}$; under this identification, $\mathcal{I}(\mathfrak{X}^n, \mathfrak{X}^m)$ is identified with $M_{m,n}(\mathcal{I}(\mathfrak{X}))$.

Note that we have equipped $M_{m,n}(\mathcal{B}(\mathfrak{X}))$ with two apparently different norms, one coming from the identification with $\mathcal{B}(\mathfrak{X}^n, \mathfrak{X}^m)$ and another arising from the embedding in $\mathcal{B}(\mathcal{B}(\mathfrak{X})^n, \mathcal{B}(\mathfrak{X})^m)$. Happily, these norms coincide, as is easily checked.

2 Elementary Results

In this section we prove that both K -groups are zero for the algebra of operators on many of the classical Banach spaces, including c_0 , $C(M)$ (M an infinite, compact metric space), ℓ_p and $L_p([0, 1])$ ($p \in [1, \infty]$).

First we consider the group K_0 . Let $\mathcal{A}$ be an algebra and define

$$\begin{aligned} \text{IP}(\mathcal{A}) &:= \{P \in \mathcal{A} \mid P^2 = P\}, \text{ the set of idempotents in } \mathcal{A}, \\ \text{IP}_n(\mathcal{A}) &:= \text{IP}(M_n(\mathcal{A})) \quad (n \in \mathbb{N}), \quad \text{IP}_\infty(\mathcal{A}) := \bigcup_{n \in \mathbb{N}} \text{IP}_n(\mathcal{A}). \end{aligned}$$

Let $m, n \in \mathbb{N}$. For $P \in \text{IP}_m(\mathcal{A})$ and $Q \in \text{IP}_n(\mathcal{A})$ we say that $P \sim_0 Q$ (in $\text{IP}_\infty(\mathcal{A})$) provided that there are matrices $R \in M_{m,n}(\mathcal{A})$ and $T \in M_{n,m}(\mathcal{A})$ satisfying: $P = RT$ and $Q = TR$. Clearly $\sim_0$ is an equivalence relation on $\text{IP}_\infty(\mathcal{A})$, and so we may form the quotient $V(\mathcal{A}) := \text{IP}_\infty(\mathcal{A})/\sim_0$. Let $[P]_V$ denote the equivalence class of $P \in \text{IP}_\infty(\mathcal{A})$. One easily checks that the operation

$$([P]_V, [Q]_V) \mapsto \left[\begin{pmatrix} P & 0 \\ 0 & Q \end{pmatrix} \right]_V, \quad V(\mathcal{A}) \times V(\mathcal{A}) \longrightarrow V(\mathcal{A}),$$

is well-defined and turns $V(\mathcal{A})$ into a commutative semigroup.

Now suppose that the algebra $\mathcal{A}$ is unital. Then we define $K_0(\mathcal{A})$ to be the Grothendieck group of $V(\mathcal{A})$ and, for $P \in \text{IP}_\infty(\mathcal{A})$, we denote the canonical image of $[P]_V$ in $K_0(\mathcal{A})$ by $[P]_0$. For $P, Q \in \text{IP}_\infty(\mathcal{A})$ we observe that

$$[P]_0 = [Q]_0 \text{ if and only if } \begin{pmatrix} P & 0 \\ 0 & I^{(k)} \end{pmatrix} \sim_0 \begin{pmatrix} Q & 0 \\ 0 & I^{(k)} \end{pmatrix} \text{ for some } k \in \mathbb{N}, \quad (2.1)$$

and we have the following *standard picture* of $K_0(\mathcal{A})$:

$$K_0(\mathcal{A}) = \{[P]_0 - [Q]_0 \mid P, Q \in \text{IP}_\infty(\mathcal{A})\}. \quad (2.2)$$

Moreover, we note that, for each $n \in \mathbb{N}$ and $P \in \text{IP}_n(\mathcal{A})$, the identity

$$[P]_0 + [I^{(n)} - P]_0 = [I^{(n)}]_0 \quad (2.3)$$

holds.

In the non-unital case, $K_0(\mathcal{A})$ is defined as a subgroup of $K_0(\mathcal{A}^\#)$ in the following way. Let $s : \mathcal{A}^\# \rightarrow \mathcal{A}^\#$ denote the *scalar map* given by $s(A + \zeta I) := \zeta I$ ($A \in \mathcal{A}$, $\zeta \in \mathbb{C}$). This is clearly an algebra homomorphism, and we define

$$K_0(\mathcal{A}) := \{[P]_0 - [s_n(P)]_0 \mid n \in \mathbb{N}, P \in \text{IP}_n(\mathcal{A}^\#)\}, \quad (2.4)$$

where we recall that s_n denotes the map on $M_n(\mathcal{A}^\#)$ induced by s .

In the case where $\mathcal{A} = \mathcal{B}(\mathfrak{X})$, the equivalence relation $\sim_0$ has a nice standard characterization.

Proposition 2.1 For $P, Q \in \text{IP}_\infty(\mathcal{B}(\mathfrak{X}))$, $P \sim_0 Q$ if and only if $\text{im } P$ is linearly homeomorphic to $\text{im } Q$. $\square$

Lemma 2.2 Suppose that $\mathfrak{X}^n$ is linearly homeomorphic to $\mathfrak{X}^{n+k}$ for some $n, k \in \mathbb{N}$. Then

$$K_0(\mathcal{B}(\mathfrak{X})) = \{[P]_0 - [Q]_0 \mid P, Q \in \text{IP}_n(\mathcal{B}(\mathfrak{X}))\}.$$

Proof. By (2.2), it suffices to prove that, for each $m \in \mathbb{N}$ and $P \in \text{IP}_m(\mathcal{B}(\mathfrak{X}))$, there is an operator $\tilde{P} \in \text{IP}_n(\mathcal{B}(\mathfrak{X}))$ with $P \sim_0 \tilde{P}$. To this end, take $j \in \mathbb{N}$ for which $n + jk \geq m$. By assumption, there are operators $R \in M_{n+jk, n}(\mathcal{B}(\mathfrak{X}))$ and $T \in M_{n, n+jk}(\mathcal{B}(\mathfrak{X}))$ satisfying $RT = I^{(n+jk)}$ and $TR = I^{(n)}$. Let $\tilde{P} := T(P \oplus 0)R$. Then clearly $\tilde{P} \in \text{IP}_n(\mathcal{B}(\mathfrak{X}))$ and $\tilde{P} \sim_0 P \oplus 0 \sim_0 P$, as desired. $\square$

A Banach space $\mathfrak{X}$ is said to be *primary* provided that, whenever $P \in \mathcal{B}(\mathfrak{X})$ is idempotent, then $\text{im } P$ or $\text{im}(I - P)$ is linearly homeomorphic to $\mathfrak{X}$.

Proposition 2.3 Suppose that $\mathfrak{X}$ is primary and linearly homeomorphic to its square $\mathfrak{X}^2$. Then $K_0(\mathcal{B}(\mathfrak{X})) = \{0\}$.

Proof. First we note that $[I]_0 = 0$ because, by assumption, $\text{im } I$ is linearly homeomorphic to $\text{im } I^{(2)}$, and so $[I]_0 = [I^{(2)}]_0 = [I]_0 + [I]_0$ (cf. Proposition 2.1).

By Lemma 2.2, it suffices to prove that $[P]_0 = 0$ for each $P \in \text{IP}(\mathcal{B}(\mathfrak{X}))$. If $\text{im } P$ is linearly homeomorphic to $\mathfrak{X}$, then $[P]_0 = [I]_0 = 0$. Otherwise, $\text{im}(I - P)$ is linearly homeomorphic to $\mathfrak{X}$, and the result follows from the identity (2.3). $\square$

Remark 2.4 As the following example shows, the condition that $\mathfrak{X}$ is linearly homeomorphic to its square cannot be removed from Proposition 2.3, not even if we require $\mathfrak{X}$ to be *prime* (i.e., the image of *every* idempotent of infinite rank is linearly homeomorphic to $\mathfrak{X}$), instead of primary.

Let $\mathfrak{X}$ denote the Banach space constructed by Gowers and Maurey in [GM2, §4.2]. This space is prime and satisfies: for $m, n \in \mathbb{N}$, $\mathfrak{X}^m$ is linearly homeomorphic to $\mathfrak{X}^n$ if and only if $m = n$. It follows immediately from this, Proposition 2.1, and (2.1) that $[I]_0$ is of infinite order in $K_0(\mathcal{B}(\mathfrak{X}))$; in particular, $K_0(\mathcal{B}(\mathfrak{X})) \neq \{0\}$.

We shall return to this example in Remark 4.4.

Remark 2.5 A natural question in relation to Proposition 2.3 is the following. Suppose that $\mathfrak{X}$ is linearly homeomorphic to its square and $K_0(\mathcal{B}(\mathfrak{X})) = \{0\}$. Is $\mathfrak{X}$ primary?

The answer to this question is ‘no’. Take $p, q \in [1, \infty[$ with $p \neq q$. Then clearly $\ell_p \oplus \ell_q$ is linearly homeomorphic to its square and $K_0(\mathcal{B}(\ell_p \oplus \ell_q)) = \{0\}$ (cf. Corollary 5.3, below), but $\ell_p \oplus \ell_q$ is not primary, because neither ℓ_p nor ℓ_q is linearly homeomorphic to $\ell_p \oplus \ell_q$.

We now proceed to consider the group K_1 . Suppose that $\mathcal{A}$ is a unital Banach algebra, and define $\text{Inv}_\infty(\mathcal{A}) := \bigcup_{n \in \mathbb{N}} \text{Inv}_n(\mathcal{A})$. Let $m, n \in \mathbb{N}$. For $U \in \text{Inv}_m(\mathcal{A})$ and $V \in \text{Inv}_n(\mathcal{A})$

we say that $U \sim_1 V$ (in $\text{Inv}_\infty(\mathcal{A})$) provided that there are a natural number $k > \max\{m, n\}$ and a continuous path $t \mapsto W_t$, $[0, 1] \rightarrow \text{Inv}_k(\mathcal{A})$, with

$$W_0 = \begin{pmatrix} U & 0 \\ 0 & I^{(k-m)} \end{pmatrix} \quad \& \quad W_1 = \begin{pmatrix} V & 0 \\ 0 & I^{(k-n)} \end{pmatrix}.$$

Clearly $\sim_1$ is an equivalence relation on $\text{Inv}_\infty(\mathcal{A})$, and so we may define

$$K_1(\mathcal{A}) := \text{Inv}_\infty(\mathcal{A}) / \sim_1.$$

Let $[U]_1$ denote the equivalence class of $U \in \text{Inv}_\infty(\mathcal{A})$. It is easily checked that the operation

$$([U]_1, [V]_1) \mapsto \left[\begin{pmatrix} U & 0 \\ 0 & V \end{pmatrix} \right]_1, \quad K_1(\mathcal{A}) \times K_1(\mathcal{A}) \longrightarrow K_1(\mathcal{A}),$$

is well-defined and turns $K_1(\mathcal{A})$ into a commutative group.

For a non-unital Banach algebra $\mathcal{A}$, we define $K_1(\mathcal{A}) := K_1(\mathcal{A}^\#)$.

Proposition 2.6 *Suppose that $\mathfrak{X}$ is linearly homeomorphic to its square $\mathfrak{X}^2$ and that the invertible group $\text{Inv}(\mathcal{B}(\mathfrak{X}))$ is connected. Then $K_1(\mathcal{B}(\mathfrak{X})) = \{0\}$.*

Proof. Let $n \in \mathbb{N}$. Clearly $\mathfrak{X}$ is linearly homeomorphic to $\mathfrak{X}^n$, and so the fact that $\text{Inv}(\mathcal{B}(\mathfrak{X}))$ is connected implies that $\text{Inv}_n(\mathcal{B}(\mathfrak{X}))$ is connected. Now the result follows from the definition of K_1 . $\square$

Remark 2.7 In fact Proposition 2.6 is just a special case of a much more general result. A unital algebra $\mathcal{A}$ is said to be *properly infinite* provided that there are elements $A_1, A_2, B_1, B_2 \in \mathcal{A}$ satisfying $B_l A_k = \delta_{kl} I$ ($k, l \in \{1, 2\}$). One easily checks that the algebra $\mathcal{B}(\mathfrak{X})$ is properly infinite in the case where $\mathfrak{X}$ is linearly homeomorphic to $\mathfrak{X}^n$ for some $n \in \{2, 3, \dots\}$. Generalizing the proof of a C^* -algebra result (cf. [Cu, Lemma 1.2] or [Rør, Proposition 3.4]) yields: for every unital, properly infinite Banach algebra $\mathcal{A}$, the group homomorphism $U \mapsto [U]_1$, $\text{Inv}(\mathcal{A}) \rightarrow K_1(\mathcal{A})$, is surjective. In particular, connectedness of $\text{Inv}(\mathcal{A})$ implies that $K_1(\mathcal{A}) = \{0\}$.

Example 2.8 (i) Let $\mathfrak{X}$ denote one of the spaces c_0 or ℓ_p ($p \in [1, \infty]$). Then certainly $\mathfrak{X}$ is linearly homeomorphic to its square and, by [LT2, Theorems 2.a.3 and 2.a.7], $\mathfrak{X}$ is prime and thus primary. Moreover, the invertible group $\text{Inv}(\mathcal{B}(\mathfrak{X}))$ is contractible, cf. [Mit, the Corollaries of Proposition 2 (p. 72) and Lemma 11b (p. 79)], and consequently connected. Hence, we obtain

$$K_\mu(\mathcal{B}(c_0)) = \{0\} \quad \& \quad K_\mu(\mathcal{B}(\ell_p)) = \{0\} \quad (p \in [1, \infty], \mu \in \{0, 1\}).$$

(ii) Let $p \in [1, \infty]$. Clearly $L_p([0, 1])$ is linearly homeomorphic to its square; it is also primary. (This is proved in [Mau] for $p = 1$ and in [AEO, Theorem 1.3] for $1 < p < \infty$;

for $p = \infty$ this follows from (i), because $L_\infty([0, 1])$ is linearly homeomorphic to ℓ_∞ , cf. [Pe].) Furthermore, by [Mit, Theorems 2a (p. 75) and 4 (p. 81)], $\text{Inv}(\mathcal{B}(L_p([0, 1])))$ is contractible and thus connected, so we conclude that

$$K_\mu(\mathcal{B}(L_p([0, 1]))) = \{0\} \quad (p \in [1, \infty], \mu \in \{0, 1\}).$$

- (iii) Let M be an infinite, compact metric space. It is well-known that $C(M)$ is linearly homeomorphic to its square. Moreover, $C(M)$ is primary with $\text{Inv}(\mathcal{B}(C(M)))$ contractible. (For M countable, this is proved in [AB, Theorems 1 and 2], and for M uncountable, this follows from [LP, Corollary 1] and [Mit, Theorem 2 (p. 73)].) Hence,

$$K_\mu(\mathcal{B}(C(M))) = \{0\} \quad (\mu \in \{0, 1\}).$$

All the results in this section (except Remark 2.4) are negative in the sense that they state that the K -groups of the algebra of operators on certain Banach spaces are trivial. To construct K -theoretically more interesting examples, we have to bring the six-term exact sequence into play. In the next section we prepare the way for this.

3 K -Theory for the Closed Ideals of Operators Contained in $\mathcal{S}$.

In this section we compute the K -groups of $\mathcal{I}(\mathfrak{X})$, where $\mathcal{I}$ is any non-zero, closed ideal of operators contained in the ideal $\mathcal{S}$ of strictly singular operators. Throughout this section, $\mathcal{I}$ denotes such an ideal of operators. The results we obtain generalize well-known results for the compact operators on a Hilbert space.

We denote the semigroup of Fredholm operators on $\mathfrak{X}$ by $\Phi(\mathfrak{X})$, and, for an operator $T \in \mathcal{B}(\mathfrak{X})$, we define the *essential spectrum* by

$$\sigma_{\text{ess}}(T) := \{\zeta \in \mathbb{C} \mid T - \zeta I \notin \Phi(\mathfrak{X})\}.$$

This is a closed, non-empty subset of the spectrum of T .

Proposition 3.1 *Let $n \in \mathbb{N}$, and let $(T_{kl})_{k,l=1}^n \in M_n(\mathcal{I}(\mathfrak{X})^\#)$. The spectrum of $(T_{kl})_{k,l=1}^n$ is independent of whether it is calculated in $M_n(\mathcal{I}(\mathfrak{X})^\#)$ or $M_n(\mathcal{B}(\mathfrak{X}))$:*

$$\sigma_{M_n(\mathcal{I}(\mathfrak{X})^\#)} \left((T_{kl})_{k,l=1}^n \right) = \sigma_{M_n(\mathcal{B}(\mathfrak{X}))} \left((T_{kl})_{k,l=1}^n \right),$$

and it is countable.

Proof. Write $T_{kl} = S_{kl} + \zeta_{kl}I$, where $S_{kl} \in \mathcal{I}(\mathfrak{X})$ and $\zeta_{kl} \in \mathbb{C}$. Since $(S_{kl})_{k,l=1}^n$ is strictly singular, it follows from [LT2, Proposition 2.c.10] that

$$\sigma_{\text{ess}} \left((T_{kl})_{k,l=1}^n \right) = \sigma_{\text{ess}} \left((\zeta_{kl}I)_{k,l=1}^n \right) \subseteq \sigma_{M_n(\mathcal{B}(\mathfrak{X}))} \left((\zeta_{kl}I)_{k,l=1}^n \right) \subseteq \sigma_{M_n(\mathbb{C})} \left((\zeta_{kl})_{k,l=1}^n \right).$$

In particular we see that $\sigma_{\text{ess}}\left((T_{kl})_{k,l=1}^n\right)$ is finite, and so, according to [Ka2, Theorem IV-(5.33)], $\sigma_{M_n(\mathcal{B}(\mathfrak{X}))}\left((T_{kl})_{k,l=1}^n\right)$ is countable. Now the result follows from [Ru, Corollary (a) of Theorem 10.18]. $\square$

For a unital Banach algebra $\mathcal{A}$, we denote the component of $\text{Inv}(\mathcal{A})$ containing the identity by $\text{Inv}^0(\mathcal{A})$.

Lemma 3.2 *Let $\mathcal{A}$ be a unital Banach algebra, and suppose that $A \in \text{Inv}(\mathcal{A})$ has countable spectrum. Then $A \in \text{Inv}^0(\mathcal{A})$.*

Proof. Take $v \in [0, \pi[$ with $\sigma(A) \cap (\mathbb{R}e^{iv}) = \emptyset$. Then $te^{iv}I + (1-t)A \in \text{Inv}(\mathcal{A})$ for all $t \in [0, 1]$, and so there is a continuous path in $\text{Inv}(\mathcal{A})$ from A to $e^{iv}I \in \text{Inv}^0(\mathcal{A})$. $\square$

Combining this with Proposition 3.1 and the definition of the group K_1 yields:

Corollary 3.3 $K_1(\mathcal{I}(\mathfrak{X})) = \{0\}$. $\square$

Proposition 3.1 has another useful consequence. Recall that, for idempotents P and Q in an algebra $\mathcal{A}$, we have $P \leq Q$ if and only if $PQ = P = QP$. The algebra $\mathcal{A}$ is said to be *finite* provided that, whenever P and Q are idempotents in $\mathcal{A}$ satisfying $P \sim_0 Q$ and $P \leq Q$, then $P = Q$. If all the algebras $M_n(\mathcal{A})$ ($n \in \mathbb{N}$) are finite, then we say that $\mathcal{A}$ is *stably finite*. Note that in the unital case, $\mathcal{A}$ is finite if and only if every left-invertible (or, equivalently, right-invertible) element in $\mathcal{A}$ is invertible.

Now suppose that $\mathcal{A}$ is a unital Banach algebra which is not finite, and take $A, B \in \mathcal{A}$ with $AB = I$ and $BA \neq I$. In particular, $A \neq 0$, and for $\zeta \in \mathbb{C}$ we have

$$|\zeta| < \frac{1}{\|A\|} \implies I - \zeta A \in \text{Inv}(\mathcal{A}) \iff A(B - \zeta I) \in \text{Inv}(\mathcal{A}) \implies \zeta \in \sigma(B).$$

Consequently, $\sigma(B)$ has non-empty interior. Combining this reasoning with Proposition 3.1 yields:

Proposition 3.4 *The Banach algebra $\mathcal{I}(X)^\sharp$ is stably finite.* $\square$

To compute $K_0(\mathcal{I}(\mathfrak{X}))$, we require the following diagonalization lemma. It is essentially just a restatement of a theorem by Edelstein and Wojtaszczyk (cf. [EW, Proposition 3.3 and Theorem 3.5] or [LT2, Theorem 2.c.13]), and indeed their proof can be taken over almost literally; only minor adjustments are necessary to ensure that all the operators in question belong to (the matrix algebras over) $\mathcal{I}(\mathfrak{X})^\sharp$.

Lemma 3.5 *Let $n \in \mathbb{N}$. For each operator $P \in \text{IP}_n(\mathcal{I}(\mathfrak{X})^\sharp)$, there are idempotents $P_1, \dots, P_n \in \mathcal{F}(\mathfrak{X})^\sharp$ and an operator $\Omega \in \text{Inv}_n(\mathcal{I}(\mathfrak{X})^\sharp)$ satisfying*

$$\text{im}(\Omega P) = \text{im}(P_1 \oplus \dots \oplus P_n).$$

In particular, the equivalence $P \sim_0 P_1 \oplus \dots \oplus P_n$ holds in $\text{IP}_\infty(\mathcal{I}(\mathfrak{X})^\sharp)$. $\square$

The following lemma is elementary.

Lemma 3.6 *Let $Q \in \text{IP}_\infty(\mathcal{F}(\mathfrak{X}))$ be of rank one, and let $n \in \mathbb{N}$. For every $P \in \text{IP}_n(\mathcal{F}(\mathfrak{X}))$, the formulas*

$$[P]_0 = (\text{rk } P) [Q]_0 \quad \& \quad [I^{(n)} - P]_0 - [I^{(n)}]_0 = -(\text{rk } P) [Q]_0$$

hold in $K_0(\mathcal{I}(\mathfrak{X})^\#)$. In particular, $[P]_0 - [s_n(P)]_0 \in \mathbb{Z} [Q]_0$ for every $P \in \text{IP}(M_n(\mathcal{F}(\mathfrak{X}))^\#)$. $\square$

Theorem 3.7 *Suppose that $Q \in \text{IP}_\infty(\mathcal{F}(\mathfrak{X}))$ is of rank one. Then the map*

$$\omega : \nu \longmapsto \nu [Q]_0, \quad \mathbb{Z} \longrightarrow K_0(\mathcal{I}(\mathfrak{X})),$$

is a group isomorphism.

Proof. Clearly ω is additive.

Assume towards a contradiction that ω is not injective. Then there is a natural number ν satisfying

$$0 = \nu [Q]_0 = \underbrace{[Q \oplus \cdots \oplus Q]_0}_\nu$$

in $K_0(\mathcal{I}(\mathfrak{X}))$ and thus in $K_0(\mathcal{I}(\mathfrak{X})^\#)$, and consequently $0 \oplus \cdots \oplus 0 \oplus I^{(k)} \sim_0 Q \oplus \cdots \oplus Q \oplus I^{(k)}$ for some $k \in \mathbb{N}$, cf. (2.1). Clearly $0 \oplus \cdots \oplus 0 \oplus I^{(k)} \leq Q \oplus \cdots \oplus Q \oplus I^{(k)}$, and so, by Proposition 3.4, $0 \oplus \cdots \oplus 0 \oplus I^{(k)} = Q \oplus \cdots \oplus Q \oplus I^{(k)}$, contradicting the fact that $Q \neq 0$. Hence, ω is injective.

To show that ω is surjective, let $g \in K_0(\mathcal{I}(\mathfrak{X}))$ be given. By (2.4), we can take $n \in \mathbb{N}$ and $P \in \text{IP}_n(\mathcal{I}(\mathfrak{X})^\#)$ so that $g = [P]_0 - [s_n(P)]_0$, and, by Lemma 3.5, we can take $P_1, \dots, P_n \in \text{IP}(\mathcal{F}(\mathfrak{X})^\#)$ for which $P \sim_0 P_1 \oplus \cdots \oplus P_n$ in $\text{IP}_\infty(\mathcal{I}(\mathfrak{X})^\#)$. Consequently,

$$g = [P_1 \oplus \cdots \oplus P_n]_0 - [s_n(P_1 \oplus \cdots \oplus P_n)]_0 = \sum_{j=1}^n ([P_j]_0 - [s(P_j)]_0) \in \mathbb{Z} [Q]_0,$$

cf. Lemma 3.6. $\square$

4 Splittings and H.I. Spaces

Let $\mathcal{A}$, $\mathcal{B}$, and $\mathcal{C}$ be algebras, and suppose that we have a short exact sequence

$$\Sigma : \quad \{0\} \longrightarrow \mathcal{A} \xrightarrow{\varphi} \mathcal{B} \xrightarrow{\psi} \mathcal{C} \longrightarrow \{0\}.$$

We say that Σ *splits algebraically* (or is *split exact*) provided that there is an algebra homomorphism $\theta : \mathcal{C} \rightarrow \mathcal{B}$ for which $\psi \circ \theta = \text{id}_{\mathcal{C}}$; such a map θ is called an *(algebraic) splitting*

homomorphism. The functor K_0 does not in general preserve short exact sequences, but it does preserve split exact sequences.

Now suppose that $\mathcal{A}$, $\mathcal{B}$, and $\mathcal{C}$ are Banach algebras and that φ and ψ are continuous. We say that Σ *splits strongly* (or is *strongly split exact*) provided that there is a continuous splitting homomorphism. The functor K_1 does not in general preserve short exact sequences, but it does preserve strongly split exact sequences.

A fundamental theorem of K -theory states that, to every short exact sequence Σ , where $\mathcal{A}$, $\mathcal{B}$, and $\mathcal{C}$ are Banach algebras and φ and ψ are continuous, we can associate the cyclic six-term exact sequence

$$\begin{array}{ccccc} K_0(\mathcal{A}) & \xrightarrow{K_0(\varphi)} & K_0(\mathcal{B}) & \xrightarrow{K_0(\psi)} & K_0(\mathcal{C}) \\ \delta_1 \uparrow & & & & \downarrow \delta_0 \\ K_1(\mathcal{C}) & \xleftarrow{K_1(\psi)} & K_1(\mathcal{B}) & \xleftarrow{K_1(\varphi)} & K_1(\mathcal{A}). \end{array}$$

The map δ_1 is called the *index map* because of its relation to the Fredholm index i , described in Proposition 4.1, below.

In this section we shall study the conditions under which the short exact sequence

$$\Sigma_{\mathfrak{X}} : \quad \{0\} \longrightarrow \mathcal{I}(\mathfrak{X}) \xrightarrow{\iota} \mathcal{B}(\mathfrak{X}) \xrightarrow{\pi} \mathcal{B}(\mathfrak{X})/\mathcal{I}(\mathfrak{X}) \longrightarrow \{0\}$$

splits. Here, as in the rest of this section, $\mathcal{I}$ denotes a closed, non-zero ideal of operators contained in $\mathcal{S}$. Note that in this case, by the results of §3, the six-term exact sequence has the following form

$$\begin{aligned} \{0\} &\longrightarrow K_1(\mathcal{B}(\mathfrak{X})) \xrightarrow{K_1(\pi)} K_1(\mathcal{B}(\mathfrak{X})/\mathcal{I}(\mathfrak{X})) \xrightarrow{\delta_1} K_0(\mathcal{I}(\mathfrak{X})) = \mathbb{Z}[Q]_0 \xrightarrow{K_0(\iota)} \dots \\ &\dots \xrightarrow{K_0(\iota)} K_0(\mathcal{B}(\mathfrak{X})) \xrightarrow{K_0(\pi)} K_0(\mathcal{B}(\mathfrak{X})/\mathcal{I}(\mathfrak{X})) \longrightarrow \{0\}, \end{aligned}$$

where Q is an idempotent of rank one.

Our first result is well-known for Hilbert spaces, and the proof of this special case is easily adapted to cover the Banach-space case.

Proposition 4.1 *For every $m \in \mathbb{N}$, the diagram*

$$\begin{array}{ccc} \Phi(\mathfrak{X}^m) & \xrightarrow{\pi_m} \text{Inv}_m(\mathcal{B}(\mathfrak{X})/\mathcal{I}(\mathfrak{X})) & \xrightarrow{[\cdot]_1} K_1(\mathcal{B}(\mathfrak{X})/\mathcal{I}(\mathfrak{X})) \\ \downarrow i & & \downarrow \delta_1 \\ \mathbb{Z} & \xrightarrow[\cong]{\omega} & K_0(\mathcal{I}(\mathfrak{X})) \end{array}$$

commutes.

□

As a consequence, we obtain:

Proposition 4.2 *The following assertions are equivalent:*

- (a) *the map $K_0(\iota) : K_0(\mathcal{I}(\mathfrak{X})) \longrightarrow K_0(\mathcal{B}(\mathfrak{X}))$ is injective;*
- (b) *the map $\delta_1 : K_1(\mathcal{B}(\mathfrak{X})/\mathcal{I}(\mathfrak{X})) \longrightarrow K_0(\mathcal{I}(\mathfrak{X}))$ is zero;*
- (c) *the map $K_1(\pi) : K_1(\mathcal{B}(\mathfrak{X})) \longrightarrow K_1(\mathcal{B}(\mathfrak{X})/\mathcal{I}(\mathfrak{X}))$ is surjective (and thus an isomorphism);*
- (d) *for each $m \in \mathbb{N}$, $\mathfrak{X}^m$ is not linearly homeomorphic to any of its proper, closed subspaces of finite codimension;*
- (e) *for each $m \in \mathbb{N}$, every Fredholm operator on $\mathfrak{X}^m$ has Fredholm index zero.*

These assertions are fulfilled in the case where the algebra $\mathcal{B}(\mathfrak{X})$ is stably finite.

Proof. The fact that (a), (b), and (c) are equivalent follows directly from the six-term exact sequence; note that, by Corollary 3.3, $K_1(\pi)$ is always injective.

The implication ‘(b) $\Rightarrow$ (e)’ is immediate from Proposition 4.1.

We prove that (d) implies (a) by contraposition. Suppose that $K_0(\iota)$ is not injective. Then there exist a number $j \in \mathbb{N}$ and an idempotent $P \in \mathcal{F}(\mathfrak{X}^j) \setminus \{0\}$ for which $[P]_0 = 0$ in $K_0(\mathcal{B}(\mathfrak{X}))$. By (2.1) and (2.3), this implies that $(I^{(j)} - P) \oplus I^{(k)} \sim_0 I^{(j)} \oplus I^{(k)}$ in $\text{IP}_\infty(\mathcal{B}(\mathfrak{X}))$ for some $k \in \mathbb{N}$ so, by Proposition 2.1, $\text{im}((I^{(j)} - P) \oplus I^{(k)})$ is linearly homeomorphic to $\text{im}(I^{(j)} \oplus I^{(k)}) = \mathfrak{X}^{j+k}$. Clearly $\text{im}((I^{(j)} - P) \oplus I^{(k)})$ is a proper, closed subspace of codimension $\text{rk } P$ in $\mathfrak{X}^{j+k}$.

The implication ‘(e) $\Rightarrow$ (d)’ is also proved contrapositively. Suppose that there exist a number $m \in \mathbb{N}$ and a proper, closed subspace $\mathfrak{W}$ of finite codimension in $\mathfrak{X}^m$ which is linearly homeomorphic to $\mathfrak{X}^m$. Take a linear homeomorphism $\tilde{T} : \mathfrak{W} \rightarrow \mathfrak{X}^m$, and let $\mathfrak{F}$ be a finite-dimensional complement of $\mathfrak{W}$ in $\mathfrak{X}^m$. Then $T := \tilde{T} \oplus 0|_{\mathfrak{F}} \in \mathcal{B}(\mathfrak{X}^m)$ is a Fredholm operator of index $i(T) = \dim \mathfrak{F} \geq 1$, and so (e) is not satisfied. Moreover, the last remark follows from this, for if $R := \tilde{T}^{-1} : \mathfrak{X}^m \rightarrow \mathfrak{W}$ is considered as an operator on $\mathfrak{X}^m$, then $TR = I^{(m)}$, but $RT \neq I^{(m)}$, so the algebra $\mathcal{B}(\mathfrak{X}^m)$ is not finite. $\square$

Corollary 4.3 (i) *Suppose that the short exact sequence $\Sigma_{\mathfrak{X}}$ splits algebraically. Then $K_0(\mathcal{B}(\mathfrak{X}))$ contains a subgroup isomorphic to $\mathbb{Z}$. In particular, $\Sigma_{\mathfrak{X}}$ never splits algebraically in the case where $K_0(\mathcal{B}(\mathfrak{X})) = \{0\}$.*

(ii) *Suppose that there exists a number $m \in \mathbb{N}$ for which $\mathfrak{X}^m$ is linearly homeomorphic to one of its proper, closed subspaces of finite codimension, or, equivalently, admits a Fredholm operator of non-zero index. Then $\Sigma_{\mathfrak{X}}$ does not split algebraically. $\square$*

Remark 4.4 It follows from (ii) that the condition in (i) is not sufficient for an algebraic splitting of the short exact sequence $\Sigma_{\mathfrak{X}}$ to exist. Indeed, consider the prime Banach space $\mathfrak{X}$ introduced in Remark 2.4. We saw there that the map

$$\nu \longmapsto \nu[I]_0, \quad \mathbb{Z} \longrightarrow K_0(\mathcal{B}(\mathfrak{X})),$$

is a group monomorphism, but $\Sigma_{\mathfrak{X}}$ does not split algebraically, because the primeness of $\mathfrak{X}$ implies that $\mathfrak{X}$ is linearly homeomorphic to each of its subspaces of finite codimension.

We do not know whether the equivalent conditions in Proposition 4.2 are sufficient for an algebraic splitting of $\Sigma_{\mathfrak{X}}$ to exist, but it seems unlikely.

The result of Corollary 4.3 might give the impression that the short exact sequence $\Sigma_{\mathfrak{X}}$ *never* splits. This is, however, not true. Recall that a Banach space $\mathfrak{X}$ is said to be *hereditarily indecomposable* or an *H.I. space* provided that no closed, infinite-dimensional subspace $\tilde{\mathfrak{X}}$ of $\mathfrak{X}$ admits an idempotent $P \in \mathcal{B}(\tilde{\mathfrak{X}})$ for which neither P nor $I|_{\tilde{\mathfrak{X}}} - P$ is of finite rank; an equivalent definition is that, for each $c > 0$ and each pair $(\mathfrak{X}_1, \mathfrak{X}_2)$ of infinite-dimensional subspaces of $\mathfrak{X}$, there are unit vectors $x_1 \in \mathfrak{X}_1$ and $x_2 \in \mathfrak{X}_2$ with $\|x_1 - x_2\| \leq c$. Gowers and Maurey's fundamental results about H.I. spaces are that they exist and that, for every H.I. space $\mathfrak{X}$, $\mathcal{B}(\mathfrak{X}) = \mathcal{S}(\mathfrak{X}) + \mathbb{C}I$ (cf. [GM1]). In particular, this implies that $\Sigma_{\mathfrak{X}}$ splits strongly for $\mathcal{I} = \mathcal{S}$. We shall now extend this result and derive its K -theoretical consequences.

For $m \in \mathbb{N}$, we denote the closed subalgebra of $M_m(\mathbb{C})$ of upper triangular matrices by $\mathcal{T}_m(\mathbb{C})$.

Proposition 4.5 *For each $m \in \mathbb{N}$, there exist a Banach space $\mathfrak{Y}$ and continuous algebra homomorphisms ψ and θ giving a strongly split exact sequence*

$$\Sigma_{\text{split}} : \quad \{0\} \longrightarrow \mathcal{S}(\mathfrak{Y}) \xrightarrow{\iota} \mathcal{B}(\mathfrak{Y}) \xrightleftharpoons[\theta]{\psi} \mathcal{T}_m(\mathbb{C}) \longrightarrow \{0\}.$$

Proof. Let $\mathfrak{Y}_1$ be an H.I. space, and take a descending chain $\mathfrak{Y}_1 \supseteq \mathfrak{Y}_2 \supseteq \cdots \supseteq \mathfrak{Y}_m$ of closed subspaces for which $\mathfrak{Y}_{j+1}$ is uncomplemented in $\mathfrak{Y}_j$ ($j \in \{1, \dots, m-1\}$). (The existence of a closed, uncomplemented subspace $\mathfrak{Y}_{j+1}$ in $\mathfrak{Y}_j$ follows from [LT1, Theorem 1], because $\mathfrak{Y}_j$ inherits the H.I. property from $\mathfrak{Y}_1$ and is consequently not linearly homeomorphic to a Hilbert space.) Let $\mathfrak{Y} := \bigoplus_{j=1}^m \mathfrak{Y}_j$, and consider an operator $T = (T_{kl})_{k,l=1}^m \in \mathcal{B}(\mathfrak{Y})$.

For $1 \leq k \leq l \leq m$, $\mathfrak{Y}_l$ is a closed subspace of the H.I. space $\mathfrak{Y}_k$ so, by [Fe], every operator $T_{kl} \in \mathcal{B}(\mathfrak{Y}_l, \mathfrak{Y}_k)$ is of the form $T_{kl} = S_{kl} + \zeta_{kl}I_{kl}$, where $S_{kl} \in \mathcal{S}(\mathfrak{Y}_l, \mathfrak{Y}_k)$, $\zeta_{kl} \in \mathbb{C}$, and I_{kl} denotes the inclusion map of $\mathfrak{Y}_l$ into $\mathfrak{Y}_k$; in particular, $I_{kk} = I|_{\mathfrak{Y}_k}$.

For $1 \leq l < k \leq m$, we have $\mathfrak{Y}_k \subseteq \mathfrak{Y}_l$, and so we may consider T_{kl} as an operator on $\mathfrak{Y}_l$ with $\text{im } T_{kl} \subseteq \mathfrak{Y}_k$. Hence, $T_{kl} = S_{kl} + \zeta_{kl}I|_{\mathfrak{Y}_l}$, where S_{kl} is strictly singular and $\zeta_{kl} \in \mathbb{C}$. Since $\mathfrak{Y}_k$ is of infinite codimension in $\mathfrak{Y}_l$, $\text{im } T_{kl}$ must be of infinite codimension in $\mathfrak{Y}_l$ as well, so $\zeta_{kl} = 0$, i.e., $T_{kl} = S_{kl} \in \mathcal{S}(\mathfrak{Y}_l, \mathfrak{Y}_k)$.

Hence, we conclude that every operator on $\mathfrak{Y}$ is of the form

$$T = \begin{pmatrix} \zeta_{11}I + S_{11} & \zeta_{12}I_{12} + S_{12} & \cdots & \zeta_{1m}I_{1m} + S_{1m} \\ S_{21} & \zeta_{22}I|_{\mathfrak{Y}_2} + S_{22} & \cdots & \zeta_{2m}I_{2m} + S_{2m} \\ \vdots & \vdots & \ddots & \vdots \\ S_{m1} & S_{m2} & \cdots & \zeta_{mm}I|_{\mathfrak{Y}_m} + S_{mm} \end{pmatrix},$$

where $\zeta_{kl} \in \mathbb{C}$ and $S_{kl} \in \mathcal{S}(\mathfrak{Y}_l, \mathfrak{Y}_k)$. The complex numbers ζ_{kl} are uniquely determined by T . Consequently, it makes sense to define a map

$$\psi : T \mapsto \begin{pmatrix} \zeta_{11} & \zeta_{12} & \cdots & \zeta_{1m} \\ 0 & \zeta_{22} & \cdots & \zeta_{2m} \\ \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & \cdots & \zeta_{mm} \end{pmatrix}, \quad \mathcal{B}(\mathfrak{Y}) \longrightarrow \mathcal{T}_m(\mathbb{C}).$$

Direct calculations show that ψ is an algebra homomorphism. By (1.1), $\ker \psi = \mathcal{S}(\mathfrak{Y})$.

Since the map

$$\theta : \begin{pmatrix} \zeta_{11} & \zeta_{12} & \cdots & \zeta_{1m} \\ 0 & \zeta_{22} & \cdots & \zeta_{2m} \\ \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & \cdots & \zeta_{mm} \end{pmatrix} \mapsto \begin{pmatrix} \zeta_{11}I & \zeta_{12}I_{12} & \cdots & \zeta_{1m}I_{1m} \\ 0 & \zeta_{22}I|_{\mathfrak{Y}_2} & \cdots & \zeta_{2m}I_{2m} \\ \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & \cdots & \zeta_{mm}I|_{\mathfrak{Y}_m} \end{pmatrix}, \quad \mathcal{T}_m(\mathbb{C}) \longrightarrow \mathcal{B}(\mathfrak{Y}),$$

is an algebra homomorphism for which $\psi \circ \theta = \text{id}_{\mathcal{T}_m(\mathbb{C})}$, we have completed the construction of the split exact sequence Σ_{split} ; θ is continuous because $\mathcal{T}_m(\mathbb{C})$ is finite-dimensional, and ψ is continuous because of the following easy result. $\square$

Lemma 4.6 *Let $\mathcal{B}$ and $\mathcal{C}$ be Banach algebras, let $\mathcal{I}$ be a closed ideal in $\mathcal{B}$, and suppose that we have a short exact sequence*

$$\{0\} \longrightarrow \mathcal{I} \xrightarrow{\iota} \mathcal{B} \xrightarrow{\psi} \mathcal{C} \longrightarrow \{0\}.$$

Then $\mathcal{C}$ is isomorphic to the quotient $\mathcal{B}/\mathcal{I}$ via the algebra isomorphism $\chi : \mathcal{B} + \mathcal{I} \mapsto \psi(\mathcal{B})$, $\mathcal{B}/\mathcal{I} \rightarrow \mathcal{C}$. Moreover, ψ is continuous if and only if χ is continuous. $\square$

Since $\mathcal{T}_m(\mathbb{C})$ is homotopy equivalent to its diagonal subalgebra $\mathbb{C}^m$, these two algebras have isomorphic K -groups, i.e., $K_0(\mathcal{T}_m(\mathbb{C})) \cong \mathbb{Z}^m$ and $K_1(\mathcal{T}_m(\mathbb{C})) = \{0\}$.

This, together with the strongly split exact sequence Σ_{split} and strong split exactness of the K -functors, gives:

Corollary 4.7 *Let $\mathfrak{Y}$ be the above Banach space. Then*

$$K_0(\mathcal{B}(\mathfrak{Y})) \cong \mathbb{Z}^{m+1} \quad \& \quad K_1(\mathcal{B}(\mathfrak{Y})) = \{0\}.$$

The generators of $K_0(\mathcal{B}(\mathfrak{Y}))$ are $[Q]_0, [P_1]_0, \dots, [P_m]_0$, where Q is an idempotent of rank one and P_j denotes the idempotent operator on $\mathfrak{Y}$ given by the $(m \times m)$ -matrix with $I|_{\mathfrak{Y}_j}$ in position (j, j) and zeroes everywhere else. $\square$

5 The Main Result

Recall that Banach spaces $\mathfrak{X}$ and $\mathfrak{Y}$ are said to be *totally incomparable* provided that no infinite-dimensional subspaces $\tilde{\mathfrak{X}}$ and $\tilde{\mathfrak{Y}}$ of $\mathfrak{X}$ and $\mathfrak{Y}$, respectively, are linearly homeomorphic (cf. [Ro]).

Example 5.1 (i) By [LT2, p. 75], any two distinct spaces in the family $\{\ell_p \mid p \in [1, \infty] \cup \{c_0\}\}$ are totally incomparable.

(ii) Suppose that $\mathfrak{X}$ has an unconditional basis and $\mathfrak{Y}$ is an H.I. space. Then $\mathfrak{X}$ and $\mathfrak{Y}$ are totally incomparable. This follows from the fact that every closed, infinite-dimensional subspace $\tilde{\mathfrak{X}}$ of $\mathfrak{X}$ contains an unconditional basic sequence (cf. [LT2, the comment just after Problem 1.d.5]), whereas $\mathfrak{Y}$ contains no unconditional basic sequence (cf. [GM1]).

We are now ready to state and prove our main theorem. For convenience, we use the convention that $\mathbb{Z}^0 := \{0\}$, the trivial group.

Theorem 5.2 *For every pair (m, n) of non-negative integers, there is a Banach space $\mathfrak{X}$ for which*

$$K_0(\mathcal{B}(\mathfrak{X})) \cong \mathbb{Z}^m \quad \& \quad K_1(\mathcal{B}(\mathfrak{X})) \cong \mathbb{Z}^n.$$

Proof. Take $\mathfrak{X}_1, \dots, \mathfrak{X}_{n+1}$ to be distinct spaces from the family $\{\ell_p \mid p \in [1, \infty] \cup \{c_0\}\}$. It follows immediately from the six-term exact sequence and Example 2.8 (i) that

$$K_0(\mathcal{B}(\mathfrak{X}_j)/\mathcal{S}(\mathfrak{X}_j)) = \{0\} \quad \& \quad K_1(\mathcal{B}(\mathfrak{X}_j)/\mathcal{S}(\mathfrak{X}_j)) \cong \mathbb{Z} \quad (j \in \{1, \dots, n+1\}).$$

If $m = 0$, let $\mathfrak{X}_{n+2} := \{0\}$. Otherwise, let $\mathfrak{X}_{n+2} := \mathfrak{Y}$, the space constructed in the proof of Proposition 4.5. Note that in both cases

$$K_0(\mathcal{B}(\mathfrak{X}_{n+2})/\mathcal{S}(\mathfrak{X}_{n+2})) \cong \mathbb{Z}^m \quad \& \quad K_1(\mathcal{B}(\mathfrak{X}_{n+2})/\mathcal{S}(\mathfrak{X}_{n+2})) = \{0\}.$$

Let $\mathfrak{X} := \bigoplus_{j=1}^{n+2} \mathfrak{X}_j$. By Example 5.1 (i)–(ii) and [Ro, Corollary 4 (b)], $\mathfrak{X}_1, \dots, \mathfrak{X}_{n+2}$ are mutually totally incomparable, and so $\mathcal{B}(\mathfrak{X}_k, \mathfrak{X}_l) = \mathcal{S}(\mathfrak{X}_k, \mathfrak{X}_l)$ ($k \neq l$). Consequently, the obvious diagonal embedding of $\bigoplus_{j=1}^{n+2} \mathcal{B}(\mathfrak{X}_j)/\mathcal{S}(\mathfrak{X}_j)$ into $\mathcal{B}(\mathfrak{X})/\mathcal{S}(\mathfrak{X})$ is in fact a homeomorphic algebra isomorphism; in particular, this implies that

$$K_\mu(\mathcal{B}(\mathfrak{X})/\mathcal{S}(\mathfrak{X})) \cong \bigoplus_{j=1}^{n+2} K_\mu(\mathcal{B}(\mathfrak{X}_j)/\mathcal{S}(\mathfrak{X}_j)) \cong \begin{cases} \mathbb{Z}^m & \text{for } \mu = 0, \\ \mathbb{Z}^{n+1} & \text{for } \mu = 1. \end{cases}$$

Combining this with the six-term exact sequence gives the following diagram

$$\begin{array}{ccccccc} \mathbb{Z} & \xrightarrow[\cong]{\omega} & K_0(\mathcal{S}(\mathfrak{X})) & \xrightarrow{K_0(\iota)} & K_0(\mathcal{B}(\mathfrak{X})) & \xrightarrow{K_0(\pi)} & K_0(\mathcal{B}(\mathfrak{X})/\mathcal{S}(\mathfrak{X})) \cong \mathbb{Z}^m \\ \sigma \uparrow & & \uparrow \delta_1 & & & & \downarrow \\ \mathbb{Z}^{n+1} & \xleftarrow[\cong]{} & K_1(\mathcal{B}(\mathfrak{X})/\mathcal{S}(\mathfrak{X})) & \xleftarrow{K_1(\pi)} & K_1(\mathcal{B}(\mathfrak{X})) & \xleftarrow{} & \{0\}, \end{array}$$

where the right-hand-side rectangle is exact, and σ denotes the induced group homomorphism, so that the left-hand-side square commutes. In fact σ is given by addition: $\sigma(\nu_1, \dots, \nu_{n+1}) = \sum_{j=1}^{n+1} \nu_j$. This follows from a straightforward calculation based on the fact that, for Fredholm operators $T_j \in \mathcal{B}(\mathfrak{X}_j)$ ($j \in \{1, \dots, n+2\}$), the operator $T_1 \oplus \dots \oplus T_{n+2} \in \mathcal{B}(\mathfrak{X})$ is a Fredholm operator of index $\sum_{j=1}^{n+2} i(T_j)$. In particular, σ is surjective, so, by exactness, $K_0(\pi)$ is injective and thus an isomorphism, i.e., $K_0(\mathcal{B}(\mathfrak{X})) \cong \mathbb{Z}^m$. Moreover, we have that $K_1(\mathcal{B}(\mathfrak{X})) \cong \text{im } K_1(\pi) = \ker \delta_1 \cong \ker \sigma \cong \mathbb{Z}^n$. $\square$

Careful tracking of the isomorphisms in the above proof shows that the generators of $K_0(\mathcal{B}(\mathfrak{X}))$ and $K_1(\mathcal{B}(\mathfrak{X}))$ are given as follows.

The group $K_0(\mathcal{B}(\mathfrak{X}))$ is generated by the elements $\underbrace{[0 \oplus \dots \oplus 0 \oplus P_j]_0}_{n+1}$ ($j \in \{1, \dots, m\}$),

where the idempotents P_j are defined as in Corollary 4.7.

The generators of $K_1(\mathcal{B}(\mathfrak{X}))$ are $[U_j]_1$ ($j \in \{1, \dots, n\}$) with the invertible operator U_j given by the matrix

$$\begin{pmatrix} I & & & & & & \\ & \ddots & & & & & \\ & & I & & & & \\ & & & L_j & & & \\ & & & & I & & \\ & & & & & \ddots & \\ & & & & & & I \\ & & A_j & & & & R \\ & & & & & & & I \end{pmatrix}.$$

Here L_j (in position (j, j)) denotes the unilateral left-shift on $\mathfrak{X}_j$, R (in position $(n+1, n+1)$) denotes the unilateral right-shift on $\mathfrak{X}_{n+1}$, and A_j (in position $(n+1, j)$) is the rank-one operator defined by $(\zeta_k)_{k \in \mathbb{N}} \mapsto (\zeta_1, 0, 0, \dots)$, $\mathfrak{X}_j \rightarrow \mathfrak{X}_{n+1}$.

As a special case of Theorem 5.2 and its proof we note that:

Corollary 5.3 *Let $n \in \mathbb{N}$, and let $\mathfrak{X}_1, \dots, \mathfrak{X}_{n+1}$ be distinct spaces belonging to the family $\{l_p \mid p \in [1, \infty] \cup \{c_0\}\}$. Then*

$$K_0\left(\mathcal{B}\left(\bigoplus_{j=1}^{n+1} \mathfrak{X}_j\right)\right) = \{0\} \quad \& \quad K_1\left(\mathcal{B}\left(\bigoplus_{j=1}^{n+1} \mathfrak{X}_j\right)\right) \cong \mathbb{Z}^n. \quad \square$$

As we have seen, every pair of torsion-free, finitely generated, commutative groups can arise as $(K_0(\mathcal{B}(\mathfrak{X})), K_1(\mathcal{B}(\mathfrak{X})))$. Our last result shows that $K_0(\mathcal{B}(\mathfrak{X}))$ may have torsion.

Proposition 5.4 *For every $k \in \{2, 3, \dots\}$, there is a Banach space $\mathfrak{X}$ for which $K_0(\mathcal{B}(\mathfrak{X}))$ contains an element of order k .*

Proof. In [GM2, §4.4], Gowers and Maurey describe how to construct a Banach space $\mathfrak{X}$ satisfying: $\mathfrak{X}^m$ is linearly homeomorphic to $\mathfrak{X}^n$ if and only if $m \equiv n \pmod{k}$. Hence, it follows from (2.1) and Proposition 2.1 that, for every $\nu \in \mathbb{N}$, $\nu [I]_0 = 0$ if and only if k divides ν . Consequently, $[I]_0$ has order k . $\square$

Acknowledgements

I want to thank N. J. Nielsen for sharing with me his insight in the theory of Banach spaces and, in particular, for introducing me to the work of Gowers and Maurey.

I should also like to thank M. Rørdam for teaching me all I know about K -theory, for many profitable discussions and, most of all, for suggesting to me to compute the K -groups of the ideal $\mathcal{S}$ of strictly singular operators. This, in fact, started the whole project.

Part of this work was carried out while I visited the University of Leeds, England, from August to December 1996. I want to thank everybody in Leeds for an excellent stay. In particular, I should like to thank H. G. Dales for his help with the Banach algebraic aspects of my work, especially for drawing my attention to the question of when the short exact sequence $\Sigma_{\mathfrak{X}}$ of §4 splits.

Finally, I want to thank the three above-mentioned persons for their careful reading of preliminary versions of this paper.

References

- [AB] D. Alspach & Y. Benyamini, Primariness of Spaces of Continuous Functions on Ordinals, *Israel J. Math.* **27** (1977), pp. 64–92.
- [AEO] D. Alspach, P. Enflo & E. Odell, On the Structure of Separable $\mathcal{L}_p$ Spaces ($1 < p < \infty$), *Studia Math.* **60** (1977), pp. 79–90.
- [Bl] B. Blackadar, *K-Theory for Operator Algebras*, M. S. R. I. Publications, Springer Verlag, 1986.
- [Cu] J. Cuntz, K -Theory for Certain C^* -Algebras, *Ann. of Math.* **113** (1981), pp. 181–197.
- [EW] I. S. Edelstein & P. Wojtaszczyk, On Projections and Unconditional Bases in Direct Sums of Banach Spaces, *Studia Math.* **56** (1976), pp. 263–276.
- [Fe] V. Ferenczi, Operators on Subspaces of Hereditarily Indecomposable Banach Spaces, *Bull. London Math. Soc.*, to appear.
- [GM1] W. T. Gowers & B. Maurey, The Unconditional Basic Sequence Problem, *J. Amer. Math. Soc.* **6** (1993), pp. 851–874.
- [GM2] W. T. Gowers & B. Maurey, Banach Spaces with Small Spaces of Operators, *preprint*.

- [Ka1] T. Kato, Perturbation Theory for Nullity Deficiency and Other Quantities of Linear Operators, *J. Analyse Math.* **6** (1958), pp. 273–322.
- [Ka2] T. Kato, *Perturbation Theory for Linear Operators*, Die Grundlehren der mathematischen Wissenschaften in Einzeldarstellungen **132**, Springer Verlag, 1966.
- [LP] J. Lindenstrauss & A. Pełczyński, Contributions to the Theory of the Classical Banach Spaces, *J. Funct. Anal.* **8** (1971), pp. 225–249.
- [LT1] J. Lindenstrauss & L. Tzafriri, On the Complemented Subspaces Problem, *Israel J. Math.* **9** (1971), pp. 263–269.
- [LT2] J. Lindenstrauss & L. Tzafriri, *Classical Banach Spaces I*, Ergebnisse der Mathematik und ihrer Grenzgebiete **92**, Springer Verlag, 1977.
- [Mau] B. Maurey, Sous-espaces complémentés de L_p , d’après P. Enflo, *Seminaire Maurey-Schwartz* 1974–1975, Ecole Polytechnique.
- [Mit] B. S. Mityagin, The Homotopy Structure of the Linear Group of a Banach Space, *Russian Mathematical Surveys* **25** (1970), pp. 59–103.
- [Peł] A. Pełczyński, On the Isomorphism Between m and M , *Bull. Acad. Pol. Sci.* **6** (1958), pp. 695–696.
- [Ro] H. P. Rosenthal, On Totally Incomparable Banach Spaces, *J. Funct. Anal.* **4** (1969), pp. 167–175.
- [Ru] W. Rudin, *Functional Analysis*, 2nd edn., McGraw-Hill, 1991.
- [Rør] M. Rørdam, Classification of Inductive Limits of Cuntz Algebras, *J. reine angew. Math.* **440** (1993), pp. 175–200.
- [Tay] J. L. Taylor, Banach Algebras and Topology, in *Algebras in Analysis* (ed. J. H. Williamson), Academic Press, 1975.